

DPNU-98-22
 hep-ph/9806273
 June 1998

A New Model of Direct Gauge Mediation with an Affine Quantum Moduli Space

Nobuhito Maru¹

*Department of Physics, Nagoya University
 Nagoya 464-8602, JAPAN*

maru@eken.phys.nagoya-u.ac.jp

Abstract

We present a new, simple supersymmetry breaking model with direct gauge mediation. Supersymmetry breaking sector is based on the model with an Affine quantum moduli space. The model has no gauge messengers and no light scalars charged under the Standard Model which give a negative contribution to the soft masses. Large expectation value at the minimum is obtained by balancing the runaway potential and the dimension six nonrenormalizable term. This makes it easy to preserve the perturbative unification. Enough suppression of the supergravity contribution to the soft masses can also be shown.

PACS: 11.30.Pb, 12.60.Jv

Keywords: Dynamical Supersymmetry Breaking, Direct Gauge Mediation,
 Affine Quantum Moduli Space

¹Research Fellow of the Japan Society for the Promotion of Science

Gauge mediated supersymmetry (SUSY) breaking [1] has been much considered as an attractive alternative to the supergravity (SUGRA) mediation since the degeneracy of the sfermions is automatically guaranteed and the soft parameters can be described in terms of a few parameters. However, the original model [1] is quite complicated, especially there is an unnatural sector called “messenger sector”. There has been much progress to simplify the structure of the model along various lines [2]-[7]. The simplest idea is “Direct Gauge Mediation”(DGM). Recently, many authors have proposed interesting DGM models [8]-[17]. For the reader who would like to know about these developments in more detail, we recommend the beautiful review of Giudice and Rattazzi [18].

In this article, we present a new, simple SUSY breaking model with direct gauge mediation.² The symmetries of our model are

$$SU(6)_1 \times SU(6)_2 \times [SU(6)], \quad (1)$$

where the first two $SU(6)_{1,2}$ are gauge groups and $SU(6)$ in the bracket is a global symmetry. We will later identify the subgroup of this $SU(6)$ with the Standard Model (SM) gauge groups.

The field content of our model is as follows.

$$\begin{aligned} X &\sim (\square, \mathbf{1}; \mathbf{1}), \\ \Sigma &\sim (\square, \bar{\square}; \mathbf{1}), \\ Q &\sim (\bar{\square}, \mathbf{1}; \square), \\ \bar{Q} &\sim (\mathbf{1}, \square; \bar{\square}). \end{aligned} \quad (2)$$

We note that this field content consists of two sectors. The first sector includes X only. This model is $SU(6) + \square$, which has been originally discussed by Csáki, Schmaltz and Skiba [19] and also discussed recently in Ref. [20, 21]. According to Ref. [19], the low energy effective theory of this model has two branches, one is $W_{dyn} = 0$ (“an Affine quantum moduli space”) and another is $W_{dyn} \neq 0$ (due to gaugino condensation). The definition of Affine quantum moduli space is that the moduli space is given by gauge invariant polynomials with no relations among them and $W_{dyn} = 0$. Moreover, ’t Hooft anomalies between fundamental fields and gauge invariant composites match at all points in the moduli space (including the origin). Therefore, if the appropriate gauge invariant operators to lift the flat directions are added to the superpotential,

²Our model has similar dynamics to the model in Ref. [16]

then we can expect the model to break SUSY due to confinement such as the model proposed by Intriligator, Seiberg and Shenker (ISS) [22]. We refer to this sector as SUSY breaking sector throughout this paper. We will utilize this branch to construct the model of direct gauge mediation. On the other hand, the case of $W_{dyn} \neq 0$, we will discuss later.

The second sector contains $\Sigma, Q, \bar{Q}$. These superfields are necessary to communicate SUSY breaking effects to the observable sector. The field content for each gauge group is $SU(6)_{1,2} + 6(\square + \bar{\square})$, which is the special case of the model in Ref. [23].

Note also that this model is completely chiral, in other words, we cannot add mass terms for any field to the superpotential.

Here we take the following tree level superpotential

$$W = \lambda_1 \Sigma Q \bar{Q} + \frac{\lambda_2}{M_P} X^4 + \frac{\lambda_3}{M_P^3} \det \Sigma, \quad (3)$$

where $\lambda_{1,2,3}$ are the couplings of order unity and M_P is the reduced Planck scale. Although it is possible to add other nonrenormalizable terms to the superpotential, we forbid them by imposing symmetries.

In the presence of the first two terms in the superpotential, there exist some classical flat directions: $v^6 \equiv \det \Sigma, B \equiv Q^6, \bar{B} \equiv \bar{Q}^6, M \equiv X Q^3$.

Let us first discuss the classical direction $\langle \Sigma \rangle \neq 0$ we are interested in, which corresponds to the direction $\det \Sigma \neq 0$;

$$\langle \Sigma \rangle = \text{diag}(v, v, v, v, v, v). \quad (4)$$

Along this direction, the gauge group $SU(6)_1 \times SU(6)_2$ is broken to their diagonal $SU(6)_D$ and $Q, \bar{Q}$ become massive since the superpotential includes the mass term $\lambda_1 \langle \Sigma \rangle Q \bar{Q}$. For large v , the low energy effective theory is $SU(6)_D + \square + \bar{\square}$ + one singlet v .³ As mentioned earlier, the low energy dynamics of this model has already been discussed [19]. $U(1)_R$ anomaly of the fermionic component of X , gaugino and that of the fermionic component of X^4 are saturated. This implies that the theory is in the confining phase below the scale Λ_L (the strong coupling scale of $SU(6) + \square + \bar{\square}$) and the low energy effective theory should be described in terms of the composite X^4 . The form of the dynamically generated superpotential due to gaugino condensation from the subgroup $SU(3) \times SU(3)$ in $SU(6)$ is

$$W_{dyn} = (\omega^r - \omega^s) \Lambda_3^3, \quad (5)$$

³We use the same notation $\langle \Sigma \rangle$ for a singlet superfield.

where ω represents the cube root of unity and Λ_3 means the strong coupling scale of pure $SU(3)$. For details of the derivation of (5), we refer the reader to Ref. [21].

What is important here is that this model has $W_{dyn} = 0$ branch. In this situation, the effective superpotential becomes

$$W_{eff} = \frac{\lambda_2}{M_P} u, \quad u \equiv X^4. \quad (6)$$

Taking into account that the effective Kähler potential is

$$K_{eff} \sim u^\dagger u / |\Lambda_L|^6, \quad (7)$$

we obtain the following scalar potential.

$$V_{eff} = \left(\frac{\partial^2 K_{eff}}{\partial u^\dagger \partial u} \right)^{-1} \left| \frac{\partial W_{eff}}{\partial u} \right|^2 = \frac{\lambda_2^2}{M_P^2} \Lambda_L^6. \quad (8)$$

At the first glance, one may think that this model is the plateau model [10, 11, 17] since the scalar potential is flat. However, we have to recall that Λ_L depends on the singlet superfield v through 1-loop matching of the gauge couplings:

$$\left(\frac{\Lambda_1}{v} \right)^9 \left(\frac{\Lambda_2}{v} \right)^{12} = \left(\frac{\Lambda_L}{\lambda_1 v} \right)^{15}, \quad (9)$$

where $\Lambda_{1,2}$ denotes the dynamical scale of $SU(6)_1 + \mathbf{\bar{6}} + 6(\mathbf{\square} + \mathbf{\bar{\square}})$ and $SU(6)_2 + 6(\mathbf{\square} + \mathbf{\bar{\square}})$, respectively. Then, the scalar potential (8) becomes

$$V_{eff} = \frac{\lambda_2^2}{M_P^2} \left(\frac{\lambda_1^5 \Lambda_1^3 \Lambda_2^4}{v^2} \right)^{\frac{6}{5}} = \frac{\lambda_2^2}{M_P^2} \left(\frac{\lambda_1^5 \Lambda^7}{v^2} \right)^{\frac{6}{5}}, \quad (10)$$

where Λ is defined as $\Lambda^7 \equiv \Lambda_1^3 \Lambda_2^4$ for simplicity. This results in runaway behavior but the term $\det \Sigma$ in Eq. (3) stabilizes the scalar potential.

Indeed, by minimizing the scalar potential

$$V_{eff} = \frac{\lambda_2^2}{M_P^2} \left(\frac{\lambda_1^5 \Lambda^7}{v^2} \right)^{\frac{6}{5}} + \frac{36 \lambda_3^2}{M_P^6} v^{10}, \quad (11)$$

we obtain

$$v \sim (\Lambda^{21} M_P^{10})^{\frac{1}{31}}, \quad F_v \sim \left(\frac{\Lambda^{105}}{M_P^{43}} \right)^{\frac{1}{31}}, \quad V_0 \sim \left(\frac{\Lambda^{210}}{M_P^{86}} \right)^{\frac{1}{31}}, \quad \frac{F_v}{v} \sim \left(\frac{\Lambda^{84}}{M_P^{53}} \right)^{\frac{1}{31}}. \quad (12)$$

Since the vacuum energy is non-zero, SUSY is certainly broken. This breaking effect is communicated to the observable sector as follows. Upon identifying the $SU(5)$

subgroup of the flavor group $SU(6)$ with our usual gauge group which includes the SM gauge groups, Q and $\bar{Q}$ behave as $\mathbf{5} + \bar{\mathbf{5}}$ messenger fields of which SUSY mass is $\lambda_1 v$ and SUSY breaking (mass)² is $\lambda_1 F_v$. They communicate SUSY breaking to the soft masses through loop diagrams of $Q, \bar{Q}$ in the usual way [1]. Note also that since the original model is completely chiral, these vectorlike messengers are dynamically generated as a result of symmetry breaking. Furthermore, there is no gauge messengers and no light scalars charged under the SM which gives a negative contribution to the soft (mass)² since Σ is a singlet for $SU(5)$.

Requiring $F_v/v \sim 10^4$ GeV to obtain the soft masses of order $10^{2\sim 3}$ GeV, we find

$$\begin{aligned}\Lambda &\sim 7 \times 10^{12} \text{GeV}, & \Lambda_L &\sim 1 \times 10^{12} \text{GeV}, \\ v &\sim 3 \times 10^{14} \text{GeV}, & \sqrt{F_v} &\sim 1 \times 10^9 \text{GeV},\end{aligned}\tag{13}$$

where we used $M_P = 2 \times 10^{18}$ GeV.

We give some comments on the above scales in order. Firstly, since $F_v < v^2$, the SM gauge groups are not broken at the minimum. Secondly, the messenger scale v is close to the GUT scale, so it is possible to preserve the perturbative unification in spite of six flavors of messengers. Thirdly, one may worry about that the SUGRA contribution to the soft masses is comparable or dominant because SUSY breaking scale $\sqrt{F_v}$ is relatively large. If we require that the gravitino mass $m_{3/2}$ which is the typical scale of the SUGRA contribution is less than 10 percent of the gluino mass,

$$m_{3/2} = \frac{F_v}{\sqrt{3}M_P} < 0.1 \times 6 \times \frac{\alpha_s}{4\pi} \frac{F_v}{v},\tag{14}$$

then we find

$$v < 0.1 \times 6 \times \frac{\alpha_s}{4\pi} \sqrt{3} M_P \sim 2 \times 10^{16} \text{GeV}.\tag{15}$$

In our model, the above requirement is clearly satisfied, so the SUGRA contribution is suppressed enough.

Recall here that there remain still classical flat directions $B, \bar{B}$ and M . We have to argue whether or not these directions are lifted quantum mechanically.

Along the B direction, $SU(6)_1$ and $SU(6)$ are completely broken. Then Σ and $\bar{Q}$ become massive, hence the low energy effective theory is $SU(6)_2 + \text{singlets}$. The dynamical superpotential is

$$W_{dyn} = \Lambda_L^3 = \Lambda_2^2 B^{1/6},\tag{16}$$

where we use the scale matching $\Lambda_L^3 = \Lambda_2^2 B^{1/6}$. This leads to non-zero constant vacuum energy. At one loop, the correction to Kähler potential makes the scalar potential stabilized near the origin [10, 11, 24].

Along the $\bar{B}$ direction, $SU(6)_2$ and $SU(6)$ are completely broken. Then Σ and Q become massive, hence the low energy effective theory is $SU(6)_1 + \square + \text{a singlet}$. The effective superpotential becomes ⁴

$$W_{eff} = \frac{\lambda_2}{M_P} u \quad (17)$$

Using the canonical Kähler potential for u and the scale matching, we obtain the scalar potential

$$V_{eff} = \frac{\Lambda_L^6}{M_P^2} = \frac{\lambda_2^2}{M_P^2} \Lambda_1^{18/5} \bar{B}^{2/5}. \quad (18)$$

Clearly, this stabilizes $\bar{B}$ direction.

Along the M direction, $SU(6)_1$ and $SU(6)$ are completely broken. The low energy effective theory is $SU(6)_2 + \text{singlets}$. This is the same effective theory along the B direction. Therefore, the M direction is also lifted.

We now briefly discuss the drawback of our model. In the case with $W_{dyn} \neq 0$, this W_{dyn} becomes a runaway potential for u, v . Then one can easily see that SUSY vacuum exists. Therefore, our model does not work in this case and it is inevitable to assume that $SU(6) + \square$ model is in $W_{dyn} = 0$ branch. The same assumption is needed if we apply the model with $\mu < \mu_{adj}$ (μ, μ_{adj} : Dynkin index of matter representation, that of the adjoint representation.) in Ref. [21] instead of $SU(6) + \square$ model. On the other hand, if we apply the model with $\mu > \mu_{adj}$ in Ref. [21], we do not have to assume $W_{dyn} = 0$ because $W_{dyn} \neq 0$ branch does not exist in this case⁵. For instance, it is an interesting challenge to construct a model with direct gauge mediation by using ISS model [22] as SUSY breaking sector.

In summary, we have presented a new, simple SUSY breaking model with direct gauge mediation. We have utilized the model with an Affine quantum moduli space as SUSY breaking sector. The model is completely chiral and has no gauge messengers and no light scalars charged under the SM gauge groups which gives a negative contribution to the soft (masses)². Since messenger fields are so heavy, the perturbative unification can be preserved. We have estimated the characteristic scales and obtain

⁴We also assume here that the theory is in the $W_{dyn} = 0$ branch.

⁵Shirman [16] uses the model that has no flat directions as SUSY breaking sector in the first example. There is also no need to assume $W_{dyn} = 0$ because it is trivial in this case.

phenomenologically desirable values. Furthermore, we have shown that the SUGRA contribution is suppressed enough.

Acknowledgements

The author would like to thank C. Csáki for valuable comments. The author also would like to thank S. Kitakado for careful reading of the manuscript. This work was supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

References

- [1] M. Dine and A.E. Nelson, Phys. Rev. **D48** (1993) 1277; M. Dine, A.E. Nelson and Y. Shirman, Phys. Rev. **D51** (1995) 1362; M. Dine, A.E. Nelson, Y. Nir and Y. Shirman, Phys. Rev. **D53** (1996) 2658.
- [2] T. Hotta, K.-I. Izawa and T. Yanagida, Phys. Rev. **D55** (1997) 415.
- [3] L. Randall, Nucl. Phys. **B495** (1997) 37.
- [4] N. Haba, N. Maru and T. Matsuoka, Nucl. Phys. **B497** (1997) 31.
- [5] N. Haba, N. Maru and T. Matsuoka, Phys. Rev. **D56** (1997) 4207.
- [6] Y. Shadmi, Phys. Lett. **B405** (1997) 99.
- [7] C. Csáki, L. Randall and W. Skiba, Phys. Rev. **D57** (1998) 383.
- [8] E. Poppits and S. Trivedi, Phys. Rev. **D55** (1997) 5508.
- [9] N. Arkani-Hamed, J. March-Russel and H. Murayama, Nucl. Phys. **B509** (1998) 3.
- [10] H. Murayama, Phys. Rev. Lett. **79** (1997) 18.
- [11] S. Dimopoulos, G. Dvali, R. Rattazzi and G. Giudice, Nucl. Phys. **B510** (1998) 12.
- [12] M.A. Luty, Phys. Lett. **B414** (1997) 71.
- [13] S. Dimopoulos, G. Dvali and R. Rattazzi, Phys. Lett. **B413** (1997) 336.

- [14] K.-I. Izawa, Y. Nomura, K. Tobe and T. Yanagida, Phys. Rev. **D56** (1997) 2886;
Y. Nomura, K. Tobe and T. Yanagida, hep-ph/9711220.
- [15] M.A. Luty and J. Terning, Phys. Rev. **D57** (1998) 6799; N. Arkani-Hamed, M.A. Luty and J. Terning, hep-ph/9712389.
- [16] Y. Shirman, Phys. Lett. **B417** (1998) 281.
- [17] K. Agashe, hep-ph/9804450.
- [18] G. Giudice and R. Rattazzi, hep-ph/9801271.
- [19] C. Csáki, M. Schmaltz and W. Skiba, Phys. Rev. **D55** (1997) 7840.
- [20] C. Csáki and H. Murayama, Nucl. Phys. **B515** (1998) 114.
- [21] G. Dotti and A.V. Manohar, Phys. Rev. Lett. **80** (1998) 2758.
- [22] K. Intriligator, N. Seiberg and S.H. Shenker, Phys. Lett. **B342** (1995) 152.
- [23] E. Poppitz, Y. Shadmi and S. Trivedi, Phys. Lett. **B388** (1996) 561.
- [24] Y. Shirman, Phys. Lett. **B389** (1996) 287.